

18 水稲うるち玄米年

銘柄名	地区別 等級	令和4年産					
		県計	内			陸	庄内
			村山	最上	置賜		
合計	構成比	100.0	28.4	12.4	17.9	58.8	41.2
	1 等	95.2	94.7	94.4	94.7	94.6	96.0
	2 等	4.0	4.6	5.1	4.5	4.7	2.9
	3 等	0.3	0.4	0.4	0.2	0.3	0.2
	規格外	0.6	0.3	0.1	0.6	0.4	0.9
はえぬき	構成比	100.0	30.1	13.0	15.6	58.7	41.3
	1 等	94.8	93.9	94.0	94.1	94.0	96.0
	2 等	4.8	5.6	5.6	5.3	5.5	3.7
	3 等	0.3	0.4	0.3	0.4	0.4	0.2
	規格外	0.1	0.1	0.1	0.1	0.1	0.0
つや姫	構成比	100.0	25.4	9.4	22.0	56.8	43.2
	1 等	98.7	98.4	99.4	97.4	98.2	99.4
	2 等	1.3	1.6	0.6	2.5	1.8	0.6
	3 等	0.0	0.0	0.0	0.0	0.0	0.0
	規格外	0.0	0.0	0.0	0.1	0.0	0.0
雪若丸	構成比	100.0	26.9	13.7	22.4	63.0	37.0
	1 等	98.0	98.4	96.8	95.7	97.1	99.5
	2 等	1.9	1.4	3.2	4.2	2.8	0.5
	3 等	0.1	0.2	0.0	0.1	0.1	0.0
	規格外	0.0	0.0	0.0	0.1	0.0	0.0
ひとめぼれ	構成比	100.0	25.9	9.1	11.4	46.4	53.6
	1 等	95.7	93.5	92.8	99.0	94.7	96.6
	2 等	4.0	6.1	6.9	0.9	5.0	3.2
	3 等	0.2	0.5	0.2	0.0	0.3	0.2
	規格外	0.0	0.0	0.1	0.0	0.0	0.0
コシヒカリ	構成比	100.0	30.1	6.6	38.6	75.3	24.7
	1 等	95.5	95.5	94.2	93.7	94.5	98.5
	2 等	4.2	3.8	5.4	6.0	5.1	1.4
	3 等	0.3	0.7	0.3	0.1	0.3	0.1
	規格外	0.1	0.1	0.0	0.2	0.1	0.0
あきたこまち	構成比	100.0	37.0	45.9	7.5	90.5	9.5
	1 等	90.2	90.7	88.9	90.8	89.8	94.3
	2 等	9.1	7.8	10.9	9.2	9.5	5.7
	3 等	0.4	1.0	0.0	0.0	0.4	0.0
	規格外	0.2	0.5	0.1	0.0	0.2	0.0
その他	構成比	100.0	19.9	10.5	22.8	53.2	46.8
	1 等	76.8	83.0	83.1	82.2	82.7	70.2
	2 等	6.8	6.1	10.0	6.2	6.9	6.6
	3 等	1.3	0.9	5.8	0.2	1.6	0.9
	規格外	15.1	10.0	1.1	11.4	8.8	22.3

- 注：1) 各年検査数量は翌3月末現在。
 2) 構成比は、検査数量の地域別割合。
 3) 四捨五入の関係で計と内訳が一致しない場合がある。
 4) 地域別の検査成績は、登録検査機関の所在する市町村に基づき集計したものである。
 (村山には県集連系で検査を行った全県分が含まれる。置賜には穀検で検査を行った全県分が含まれる。)

産別銘柄別検査成績

単位：%

銘柄名	令和3年産						銘柄名	令和2年産					
	県計	内陸				庄内		県計	内陸				庄内
		村山	最上	置賜	計				村山	最上	置賜	計	
合	100.0	27.3	13.1	19.0	59.4	40.6	合	100.0	27.4	12.7	19.9	60.1	39.9
	93.9	92.8	91.0	93.4	92.6	96.0		94.4	94.3	92.0	93.8	93.6	95.6
	5.1	6.2	8.5	5.6	6.5	3.0		4.7	4.9	7.6	5.2	5.6	3.4
	0.4	0.6	0.4	0.3	0.5	0.2		0.3	0.4	0.3	0.3	0.4	0.2
計	0.6	0.5	0.1	0.7	0.5	0.8	計	0.6	0.4	0.1	0.7	0.4	0.9
はえぬき	100.0	28.6	14.1	17.2	59.9	40.1	はえぬき	100.0	28.7	13.7	18.2	60.6	39.4
	93.8	92.6	90.6	92.8	92.2	96.2		94.6	94.3	92.6	93.3	93.6	96.0
	5.6	6.6	8.8	6.2	7.0	3.5		5.0	5.1	7.1	5.9	5.8	3.8
	0.4	0.6	0.5	0.5	0.5	0.3		0.3	0.4	0.2	0.3	0.3	0.2
	0.2	0.2	0.1	0.6	0.3	0.0		0.2	0.2	0.1	0.5	0.2	0.0
つや姫	100.0	24.9	9.7	22.4	57.0	43.0	つや姫	100.0	24.5	9.4	23.6	57.5	42.5
	97.2	95.7	96.7	96.2	96.0	98.6		98.9	98.7	99.1	97.8	98.4	99.6
	2.5	4.0	3.3	3.0	3.5	1.2		0.8	1.3	0.8	1.2	1.2	0.4
	0.1	0.3	0.0	0.1	0.1	0.1		0.0	0.0	0.0	0.0	0.0	0.0
	0.2	0.0	0.0	0.8	0.3	0.0		0.2	0.0	0.1	0.9	0.4	0.0
雪若丸	100.0	24.2	13.5	24.2	61.9	38.1	雪若丸	100.0	24.8	13.5	25.8	64.1	35.9
	97.4	96.8	94.5	96.0	96.0	99.6		97.7	97.6	94.6	96.9	96.7	99.6
	2.5	3.2	5.5	3.6	3.9	0.4		2.2	2.3	5.2	2.9	3.1	0.4
	0.0	0.0	0.0	0.1	0.0	0.0		0.0	0.1	0.1	0.0	0.1	0.0
	0.1	0.0	0.0	0.3	0.1	0.0		0.1	0.0	0.2	0.2	0.1	0.0
ひとめぼれ	100.0	23.7	9.5	10.9	44.1	55.9	ひとめぼれ	100.0	23.6	8.9	11.0	43.5	56.5
	95.4	91.9	89.4	95.8	92.3	97.8		95.1	92.4	89.3	93.3	92.0	97.6
	4.1	6.6	9.8	3.8	6.6	2.1		4.5	6.7	9.0	6.5	7.1	2.4
	0.3	1.1	0.7	0.1	0.8	0.0		0.3	0.7	1.7	0.0	0.7	0.0
	0.2	0.3	0.2	0.3	0.3	0.1		0.1	0.1	0.0	0.2	0.1	0.0
コシヒカリ	100.0	27.7	7.0	42.5	77.2	22.8	コシヒカリ	100.0	25.6	6.6	42.9	75.1	24.9
	95.4	95.2	94.5	94.0	94.5	98.5		90.4	89.3	91.2	89.3	89.5	93.1
	4.0	4.3	5.2	5.1	4.8	1.2		8.8	9.7	8.4	9.9	9.7	6.2
	0.3	0.5	0.3	0.2	0.3	0.3		0.4	0.9	0.1	0.1	0.4	0.5
	0.3	0.0	0.0	0.7	0.4	0.0		0.4	0.1	0.2	0.7	0.5	0.1
あきたこまち	100.0	37.9	46.9	6.6	91.4	8.6	あきたこまち	100.0	39.2	45.2	7.9	92.3	7.7
	83.9	84.0	81.2	87.7	82.8	95.4		79.6	85.0	71.7	82.0	78.2	96.4
	15.0	14.2	18.0	11.4	16.0	4.6		19.1	12.9	27.6	15.8	20.4	3.6
	0.8	1.3	0.5	0.2	0.8	0.0		0.8	1.2	0.5	1.2	0.9	0.0
	0.3	0.5	0.2	0.7	0.4	0.0		0.5	0.8	0.2	1.0	0.5	0.0
その他	100.0	26.4	11.0	20.4	57.8	42.2	その他	100.0	28.1	7.9	20.8	56.8	43.2
	73.5	78.9	82.8	76.6	78.8	66.2		73.3	81.2	83.9	83.3	82.4	61.4
	13.0	10.3	15.6	18.2	14.1	11.6		13.0	10.2	13.7	9.8	10.5	16.2
	1.1	1.7	1.2	0.5	1.2	0.9		1.7	1.8	0.8	3.1	2.1	1.0
	12.4	9.1	0.4	4.8	5.9	21.2		12.1	6.8	1.6	3.8	5.0	21.4

(山形県農業技術環境課調べ)